



## Modelling and computation for nanoparticle reconstruction using Electron Energy Loss Spectroscopy

Karamehmedovi, Mirza; Kiewidt, Lars; Wriedt, Thomas

*Published in:*  
Book of Abstracts, META'14

*Publication date:*  
2014

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Karamehmedovi, M., Kiewidt, L., & Wriedt, T. (2014). Modelling and computation for nanoparticle reconstruction using Electron Energy Loss Spectroscopy. In S. Zouhdi, N. Zheludev, & J. H. Teng (Eds.), *Book of Abstracts, META'14* (pp. 1182)  
<http://metaconferences.org/ocs/index.php/META14/index/pages/view/proceedings/#.VKumSHu9Hx5>

---

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

# Modelling and computation for nanoparticle reconstruction using Electron Energy Loss Spectroscopy

Mirza Karamehmedović<sup>1\*</sup>, Lars Kiewidt<sup>2</sup>, and Thomas Wriedt<sup>3</sup>

<sup>1</sup>DTU Compute and DTU Physics, Technical University of Denmark, Kgs. Lyngby, Denmark

<sup>2</sup>Verfahrenstechnik der Wertstoffrückgewinnung (VdW), University of Bremen, Bremen, Germany

<sup>3</sup>Institut für Werkstofftechnik (IWT), Bremen, Germany

\*corresponding author, E-mail: mika@dtu.dk

## Abstract

In Electron Energy Loss Spectroscopy, or EELS, one measures the interaction between a nanoparticle and a probing relativistic electron beam. When the beam is steered to a small but finite distance from the particle surface, the plasmons induced in the particle emit an electromagnetic field that causes a measurable change in the kinetic energy of the beam electrons [1, 2, 3, 4, 5]. The measured electron energy loss spectrum can be used to infer the size, shape and material properties of the nanoparticle [6]. A fast and stable solution of this inverse problem requires a sparse but physically well-justified forward model. In this talk, we shall describe and discuss important issues that arise in the mathematical modelling and the numerical implementation of relevant forward models based on discrete sources. We shall pay special attention to a Generalized Multipole Technique (GMT, [7, 8]) and to the Null-Field Method with Discrete Sources (NFM-DS) of T-Matrix type [9, 10]. The models are constructed around the central equation

$$P(\omega) = \frac{e}{\pi \hbar \omega} \int_{t=-\infty}^{\infty} \Re [\mathbf{v} \cdot \mathbf{E}^{\text{sca}}(\mathbf{r}_e(t), \omega) e^{-i\omega t}] \quad (1)$$

that links the probabilistic and the classical aspects of the EELS setup, and describes the electron energy loss probability density  $P$  at energy  $\hbar\omega$  in terms of the electron beam velocity  $\mathbf{v}$  and path  $\mathbf{r}_e(t)$ , and the near field  $\mathbf{E}^{\text{sca}}$  induced by plasmons in the nanoparticle. We shall support our discussion using concrete examples and numerical results.

## References

- [1] F.J. García de Abajo, Optical excitations in electron microscopy, *Reviews of Modern Physics* 82: 209275, 2010.
- [2] G. Ruthermann, Diskrete Energieverluste mittelschneller Elektronen beim Durchgang durch dünne Folien, *Annals of Physics* 2: 113134, 1948.
- [3] H. Watanabe, Experimental evidence for the collective nature of the characteristic energy loss of electrons in solids studies on the dispersion relation of plasma frequency, *Journal of the Physical Society of Japan* 11: 112119, 1956.

- [4] C.J. Powell, J.B. Swan, Origin of the characteristic electron energy losses in aluminum, *Physical Review* 115: 869875, 1959.
- [5] H. Raether, Excitation of Plasmons and Interband Transitions by Electrons, of series Springer Tracks in Modern Physics, Springer-Verlag, Berlin, 1980 vol. 88.
- [6] R. Erni, S. Lazar, N.D. Browning, Prospects for analyzing the electronic properties in nanoscale systems by VEELS, *Ultramicroscopy* 108 270276, 2008.
- [7] L. Kiewidt, M. Karamehmedović, C. Matyssek, W. Hergert, L. Mädler, T. Wriedt, Numerical simulation of Electron Energy Loss Spectroscopy using a Generalized Multipole Technique, *Ultramicroscopy* 133: 101–108, 2013.
- [8] T. Wriedt (Ed.), Generalized Multipole Techniques for Electromagnetic and Light Scattering, Elsevier, Amsterdam, NLD, 1999.
- [9] A. Doicu, T. Wriedt and Yu. Eremin, Light Scattering by Systems of Particles, Null-Field Method with Discrete Sources: Theory and Programs (Springer 2006).
- [10] A. Doicu and T. Wriedt, Near-field computation using the null-field method, *J. Quant. Spectrosc. Radiat. Transf.* 111(3), 466473 (2010).